

FLOCCULATION OF MICROALGAE AND COHESIVE SEDIMENT


Dr. Nguyen Ho Que

Energy Environment Institute, Kyungpook National University, 2559 Gyeongsang-daero, Sangju, Gyeongbuk 37224, Korea

Faculty of Environment and Labour Safety, Ton Duc Thang University, Ho Chi Minh, Vietnam

E-mail: hoqueyt@gmail.com

Interaction between microalgae and clay minerals enhances biologically mediated flocculation, therefore affecting the sedimentation and transportation of suspended particulate matter (SPM) in the aquatic system. This interaction forms larger flocs with a higher settling velocity and controls the fate of SPM. In this research, we investigated the flocculation kinetics of microalgae and cohesive sediment to elucidate the associated mechanism in such kinds of interactions. The procedure of research is disrobed in below figure:



A flocculation jar test was applied to various mixtures of kaolinite and microalgae samples from batch culture (*Chlorella vulgaris*) to estimate the biologically mediated flocculation kinetics. The organic matter (OM) composition, secreted from the microalgae, was characterized by a liquid chromatography-organic carbon detection system, and quantitative analysis on transparent exopolymer particles was led separately. A two-class flocculation kinetic model, based on the interaction between flocculi and flocs, was also adopted to quantitatively analyze the experimental data from the flocculation. Results from the flocculation kinetic tests and OM analyses, in association with other data analyses (i.e., floc size distribution, flocculation kinetic model), showed that flocculation increased with OM concentration during the growth phase. However, on the day of the early stationary phase, flocculation kinetics started decreasing and substantially declined as time went even though the amount of OM was still increasing. Our results indicate that an adequate quantity of biopolymers produced by the microalgae cells in the growth phase enhanced floc-to-floc attachment, and hence flocculation kinetics. In contrast, an excessive quantity of biopolymers and humic substances in the stationary phase enhanced the formation of polymeric backbone structures and flocculation via scavenging particles, but it simultaneously increased steric stabilization with the production of many fragmented particles.

Special Seminar



Dr. NGUYEN HO QUE

Title: Flocculation of Microalgae and Cohesive sediment
Speaker: Dr. NGUYEN HO QUE (Ph.D. in Material Science)

Lecturer at Faculty of Environment and Labour Safety,
Ton Duc Thang University, Vietnam
Postdoctoral Research Associate in Water Quality and
Resources, Kyungpook National University, S. Korea

Date and Time: December 12, 2022, at 10:00 am

Location: 306 Meeting Room, 3F in Building of Research &
Project 1

Organized by
Applied Ecological Engineering Lab.(Prof. Fujino),
Department of Environmental Science & Technology, Saitama University

Supported by
The Strategic Research Area for Sustainable Development in East Asia (SRASDEA)

Contact: fujino@mail.satiama-u.ac.jp